



NATURA 2000 - STANDARD DATA FORM

For Special Protection Areas (SPA),
Proposed Sites for Community Importance (pSCI),
Sites of Community Importance (SCI) and
for Special Areas of Conservation (SAC)

SITE ES0000170
SITENAME Área esteparia de La Mancha norte

TABLE OF CONTENTS

- [1. SITE IDENTIFICATION](#)
- [2. SITE LOCATION](#)
- [3. ECOLOGICAL INFORMATION](#)
- [4. SITE DESCRIPTION](#)
- [5. SITE PROTECTION STATUS AND RELATION WITH CORINE BIOTOPES](#)
- [6. IMPACTS AND ACTIVITIES IN AND AROUND THE SITE](#)
- [7. MAP OF THE SITE](#)

1. SITE IDENTIFICATION

| | | |
|----------------------|-----------------------------------|-----------------------------|
| 1.1 Type A | 1.2 Site code ES0000170 | Back to top |
|----------------------|-----------------------------------|-----------------------------|

1.3 Site name

Área esteparia de La Mancha norte

| | |
|--|-----------------------------------|
| 1.4 First Compilation date 1997-12 | 1.5 Update date 2004-08 |
|--|-----------------------------------|

1.6 Respondent:

Name/Organisation: Junta de Comunidades de Castilla-La Mancha - Consejería de Agricultura - Dirección General de Montes y Espacios Naturales
Address: C/Quintanar de la Orden s/n. 45071 TOLEDO (ESPAÑA)
Email: rednaturaclm@jccm.es

1.7 Site indication and designation / classification dates

| | |
|--|---------|
| Date site classified as SPA: | 2005-07 |
| National legal reference of SPA designation | No data |

2. SITE LOCATION

2.1 Site-centre location [decimal degrees]:

[Back to top](#)

Longitude
-3.578888889

Latitude
39.68805556

2.2 Area [ha]:
107245.96

2.3 Marine area [%]
0.0

2.4 Sitelength [km]:
0.0

2.5 Administrative region code and name

NUTS level 2 code **Region Name**

| | |
|------|--------------------|
| ES42 | Castilla-La Mancha |
| ES42 | Castilla-La Mancha |

2.6 Biogeographical Region(s)

Mediterranean (0.0
%)

3. ECOLOGICAL INFORMATION

3.1 Habitat types present on the site and assessment for them

[Back to top](#)

| Annex I Habitat types | | | | | | Site assessment | | | |
|-----------------------|----|----|------------|---------------|--------------|------------------|------------------|--------------|--------|
| Code | PF | NP | Cover [ha] | Cave [number] | Data quality | A B C D | A B C | | |
| | | | | | | Representativity | Relative Surface | Conservation | Global |
| 1410 | | | | | | B | | B | |
| 1520 | | | | | | B | | B | |
| 4090 | | | | | | B | | B | |
| 5210 | | | | | | B | | B | |
| 6220 | | | | | | B | | B | |
| 92A0 | | | | | | C | | B | |
| 9340 | | | | | | B | | B | |

- **PF:** for the habitat types that can have a non-priority as well as a priority form (6210, 7130, 9430) enter "X" in the column PF to indicate the priority form.
- **NP:** in case that a habitat type no longer exists in the site enter: x (optional)
- **Cover:** decimal values can be entered

- **Caves:** for habitat types 8310, 8330 (caves) enter the number of caves if estimated surface is not available.
- **Data quality:** G = 'Good' (e.g. based on surveys); M = 'Moderate' (e.g. based on partial data with some extrapolation); P = 'Poor' (e.g. rough estimation)

3.2 Species referred to in Article 4 of Directive 2009/147/EC and listed in Annex II of Directive 92/43/EEC and site evaluation for them

| Species | | | | | Population in the site | | | | | | Site assessment | |
|---------|------|--|---|----|------------------------|------|------|------|---------|--------------|-----------------|-------|
| Group | Code | Scientific Name | S | NP | Type | Size | | Unit | Cat. | Data quality | A B C D | A B C |
| | | | | | | Min | Max | | C R V P | | Pop. | Cons. |
| B | A133 | Burhinus oedicanus | | | p | | | | C | | | B |
| B | A133 | Burhinus oedicanus | | | r | | | | C | | | B |
| B | A431 | Calandrella rufescens | | | r | | | | R | | | |
| B | A082 | Circus cyaneus | | | p | | | | P | | | B |
| B | A082 | Circus cyaneus | | | r | | | | R | | | B |
| B | A082 | Circus cyaneus | | | w | | | | R | | | B |
| B | A084 | Circus pygargus | | | r | 5 | 5 | p | P | G | | B |
| B | A084 | Circus pygargus | | | p | 5 | 5 | p | | G | | B |
| B | A231 | Coracias garrulus | | | | | | | | | | |
| B | A095 | Falco naumanni | | | r | 214 | 214 | p | | G | | B |
| B | A127 | Grus grus | | | w | 1000 | | i | | M | | |
| B | A242 | Melanocorypha calandra | | | p | | | | C | | | B |
| B | A129 | Otis tarda | | | p | 1700 | 1700 | i | | G | | A |
| B | A205 | Pterocles alchata | | | r | 2624 | 2624 | i | C | G | | B |
| B | A205 | Pterocles alchata | | | p | 2624 | 2624 | i | | G | | B |
| B | A420 | Pterocles orientalis | | | r | | | | C | | | B |
| B | A420 | Pterocles orientalis | | | p | | | | V | | | B |
| B | A128 | Tetrax tetrax | | | p | 6135 | 6135 | i | | G | | B |

- **Group:** A = Amphibians, B = Birds, F = Fish, I = Invertebrates, M = Mammals, P = Plants, R = Reptiles
- **S:** in case that the data on species are sensitive and therefore have to be blocked for any public access enter: yes
- **NP:** in case that a species is no longer present in the site enter: x (optional)
- **Type:** p = permanent, r = reproducing, c = concentration, w = wintering (for plant and non-migratory)

- species use permanent)
- **Unit:** i = individuals, p = pairs or other units according to the Standard list of population units and codes in accordance with Article 12 and 17 reporting (see [reference portal](#))
- **Abundance categories (Cat.):** C = common, R = rare, V = very rare, P = present - to fill if data are deficient (DD) or in addition to population size information
- **Data quality:** G = 'Good' (e.g. based on surveys); M = 'Moderate' (e.g. based on partial data with some extrapolation); P = 'Poor' (e.g. rough estimation); VP = 'Very poor' (use this category only, if not even a rough estimation of the population size can be made, in this case the fields for population size can remain empty, but the field "Abundance categories" has to be filled in)

4. SITE DESCRIPTION

[Back to top](#)

4.1 General site character

| Habitat class | % Cover |
|---------------------|---------|
| Total Habitat Cover | 0 |

Other Site Characteristics

Conjunto de 9 áreas de importancia para las aves esteparias, sobre todo la avutarda (Otis tarda), donde en la zona se concentra más del 60 % de la población de avutardas de Castilla-La Mancha. De relieve llano o suavemente ondulado, la zona se ubica en la comarca natural de La Mancha, entre las provincias de Cuenca, Ciudad Real y Toledo.

4.2 Quality and importance

Conjunto de áreas cultivadas con pastizal e intercalados con enclaves de encinar, coscojar y formaciones propias de estepas yesosas. La importancia de la zona radica en la cuantía de las poblaciones de aves esteparias que alberga, sobre todo de avutarda.

4.3 Threats, pressures and activities with impacts on the site

The most important impacts and activities with high effect on the site

| Negative Impacts | | | |
|------------------|------------------------------|-----------------------------|------------------------|
| Rank | Threats and pressures [code] | Pollution (optional) [code] | inside/outside [i o b] |
| H | A10 | | I |
| M | J01 | | I |
| H | A02 | | I |
| M | A09 | | O |
| M | D02.01 | | I |
| M | D04.02 | | I |
| H | A09 | | I |
| M | A07 | | I |
| H | A04.03 | | I |
| M | D01.02 | | I |

| Positive Impacts | | | |
|------------------|-------------------------------|-----------------------------|------------------------|
| Rank | Activities, management [code] | Pollution (optional) [code] | inside/outside [i o b] |
| H | A09 | | I |

Rank: H = high, M = medium, L = low

Pollution: N = Nitrogen input, P = Phosphor/Phosphate input, A = Acid input/acidification,

T = toxic inorganic chemicals, O = toxic organic chemicals, X = Mixed pollutions

i = inside, o = outside, b = both

4.4 Ownership (optional)

| Type | [%] |
|------------------|-----|
| National/Federal | 0 |

| | | |
|-----------------------|-----------------|-------------------|
| Public | State/Province | 0 |
| | Local/Municipal | 0 |
| | Any Public | 0.02 |
| Joint or Co-Ownership | | 0.27 |
| Private | | 0 |
| Unknown | | 99.71 |
| sum | | 99.99999999999999 |

4.5 Documentation

- ESPARVEL, 2002. Campaña de Conservación de la población de Aguilucho Cenizo y Pálido en Toledo, 2002. - ESPARVEL, 1997. Campaña de Conservación del aguilucho cenizo en la provincia de Toledo. Temporada de cría 1997. - ESPARVEL, 1997. Distribución y censo del aguilucho cenizo en la provincia de Toledo. Temporada de cría de 1997. - ESPARVEL, 1997. Estudio y conservación de la población del Cernícalo primilla en Toledo. - Velasco, T. 2002. Censo invernal de aves acuáticas en las zonas húmedas de la provincia de Toledo, Enero 2001, Enero 2002. - Polo, J. y Pérez, J. 2003. Resultado de los censos de aves esteparias, abril-mayo 2000, enero 2001, invierno-primavera 2001, invierno 2002 en Toledo. - TRAGSA, 2002. Censo y seguimiento anual de las poblaciones de aves esteparias de la provincia de Ciudad Real. - Viada, C. (ed.). 1998. Áreas importantes para las aves de España. 2ª Edición Revisada y Ampliada; Monografía nº 5; SEO-Birdlife.

5. SITE PROTECTION STATUS (optional)

[Back to top](#)

5.1 Designation types at national and regional level:

| Code | Cover [%] | Code | Cover [%] | Code | Cover [%] |
|------|-----------|------|-----------|------|-----------|
| ES00 | 100.0 | | | | |

6. SITE MANAGEMENT

[Back to top](#)

6.1 Body(ies) responsible for the site management:

| | |
|---------------|---|
| Organisation: | Junta de Comunidades de Castilla-La Mancha - Consejería de Agricultura - Dirección General de Montes y Espacios Naturales |
| Address: | C/Quintanar de la Orden s/n. 45071 TOLEDO (ESPAÑA) |
| Email: | rednaturaclm@jccm.es |

6.2 Management Plan(s):

An actual management plan does exist:

| | |
|-------------------------------------|------------------------|
| <input checked="" type="checkbox"/> | Yes |
| <input type="checkbox"/> | No, but in preparation |
| <input type="checkbox"/> | No |

6.3 Conservation measures (optional)

En diversos polígonos catastrales de los municipios de Fuente de Pedro Naharro, Horcajo de Santiago, Torrubia del Campo, Uclés, Almendros, Villamayor de Santiago, Pozorrubio y Horcajo de Santiago en Cuenca y de Consuegra, Corral de Almaguer, Dosbarrios, El Romeral, La Guardia, Madrudejos, Sta.Cruz de la Zarza, Tembleque, Turleque, Villacañas, Villanueva de Bogas, Villarrubia de Santiago y Villatobas en Toledo, se aplica el Programa de Ayudas a la adopción de prácticas agrícolas compatibles con la conservación del hábitat para las aves esteparias en la Comunidad Autónoma de Castilla-La Mancha (Orden de la Consejería de Agricultura y Medio Ambiente, de 24 de marzo de 1995)

7. MAP OF THE SITES

[Back to top](#)

INSPIRE ID:

Map delivered as PDF in electronic format (optional)

Yes No

Reference(s) to the original map used for the digitalisation of the electronic boundaries (optional).